

ANC an- kun²

anjir of G 0016, Jiangsu, China;
anjir of G 70, China

Pb 630.6

3 1671- 2552 20

T, activity on the n
na.

J-
liz
m
he
v
i.
an

ite and granodior
ectively. Zircon H
with possible invc
activities in Tarir
neous activities an
rica orogeny. Fro
a passive contine

f isotope; tectoni

2013- 12- 02

2014- 03- 19

41172175

1968-

mail: zchu

Ma 65
5M

Ma

2009K P
15-58% 12-15

200

100

P- MS

U-

Hf

U- Pb

U- Pb

Hf

2

o

2010KR015
150-250µ m
CL

2-3m

3

%

2%

10%-

3via 100-0.38

a

ri... c... rian strata in
nd... rocks with s
ol... eoproterozo

ordia dia... m...
d granod...

						40		
						21		
						71	0	
			1.02	0.1025	0.39	0.86		
21	600	578	1.04	0.1028	0.46	0.87		
22	4081	2256	1.81	0.1027	0.45	0.86		
23	937	989	0.95	0.0975	0.46	0.80		
24	510	494	1.03	0.1029	0.49	0.80	1.0	631

Hf

2

—

—

—

[4- 518]

[41418]

Hf

[10

612

6

4.2

Zhu

U - Pb

hu [2]

68

1- c [11,23]

Xu [11]

SH

Pb

740Ma 725Ma

-600

Zhu [7]

Pb

760Ma

820-760Ma

Li [13,24]

R

[13,24]

[13]

53

630

[5]Zha

[J].

2007, 23: 185-191.

John B M, Zhang C

Crustal growth in

1. Isotopic character

2000, 328: 15- 51.

Yuan C, Sun M, et

2007, 152: 1-10.

Neoproterozoic tectonics

Aksu blueschists in northwestern

A- ICFD and

in

[J].

Figura

Cambrian

crustal structure,

Review on Geophysics;

con U- Pb and Hf isotope

with the Emeishan mantle

2008, 72: 3084- 3104.

Neoproterozoic granitoids in South

111-118.